

Suppressing FCNC and CP-Violating Phases by Extra Dimensions

Jisuke Kubo and Haruhiko Terao

Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan

Abstract

In extra dimensions the infrared attractive force of gauge interactions is amplified. We find that this force can align in the infrared limit the soft-supersymmetry breaking terms out of their anarchical disorder at a fundamental scale, in such a way that flavor-changing neutral currents as well as dangerous CP-violating phases are sufficiently suppressed at the unification scale. The main assumption is that the matter and Higgs supermultiplets and the flavor-dependent interactions such as Yukawa interactions are stuck at the four-dimensional boundary. As a concrete example we consider the minimal model based on $SU(5)$ in six dimensions.

11.10.Hi, 11.10.Kk, 12.10.-g, 12.60.Jv

Low-energy softly-broken supersymmetry (SUSY) has been the most promising idea in solving the gauge hierarchy problem [1]. However, the introduction of the superpartners of the known particles induces large flavor-changing neutral current (FCNC) processes and CP-violating phases, which are severely constrained by precision experiments [2]. Therefore, the huge degrees of freedom involved in the soft-supersymmetry breaking (SSB) parameters have to be highly constrained in all viable supersymmetric models. This has been called the supersymmetric flavor problem.

To overcome this problem, several ideas of SUSY breaking and its mediation mechanisms have been proposed; gauge mediation [3], anomaly mediation [4], gaugino mediation [5] and so on. The common feature behind these ideas is that the SSB parameters are generated by flavor-blind radiative corrections, and that their tree-level contributions at a fundamental scale M_{PL} are assumed to be sufficiently suppressed, *e.g.*, by sequestering of branes for the visible sector and the hidden SSB sector. However, it has been argued recently [6] that such a sequestering mechanism cannot be simply realized in generic supergravity or superstring inspired models. Therefore, the supersymmetric flavor problem still manifests itself in the above-mentioned mediation mechanisms and their modifications.

An interesting way out from this problem is to suppress the tree-level contributions by certain field theoretical dynamics. There have been indeed several attempts along the line of thought, in which use has been made [7–9] that the SSB parameters are suppressed in the infrared limit in approximate superconformal field theories [10]. In this letter, we propose another possibility in more than four dimensions that flavor-blind radiative corrections dominate over the tree-level contributions as well as any other flavor non-universal corrections.

Let us start to present our idea. We assume that only the supersymmetric gauge interactions exist in the $(4 + \delta)$ dimensional bulk ($\delta = 1$ or 2), while all the other interactions are confined at the four-dimensional boundary [11–13]. Accordingly, the $(4 + \delta)$ -dimensional gauge supermultiplet propagates in the bulk, and all the $N = 1$ chiral supermultiplets $\Phi_i = (\phi_i, \psi_i)$ containing matters and Higgses propagate only in four dimensions. The gauge supermultiplet contains a chiral supermultiplet Γ in the adjoint representation, to which we assign an odd parity [12,13] so that it does not contain zero modes, and does not have any interactions with Φ 's. To simplify the situation, we further assume that each extra dimension is compactified on a circle with the same radius R . The size of R is model dependent, but throughout this letter we assume that $M_{\text{GUT}} = 1/R$. The boundary superpotential has a generic form

$$W(\Phi) = \frac{1}{6} Y^{ijk} \Phi_i \Phi_j \Phi_k + \frac{1}{2} \mu^{ij} \Phi_i \Phi_j, \quad (1)$$

and that the SSB Lagrangian L_{SSB} can be written as

$$-L_{\text{SSB}} = \left(\frac{1}{6} h^{ijk} \phi_i \phi_j \phi_k + \frac{1}{2} B^{ij} \phi_i \phi_j + \frac{1}{2} \sum_{n=0} M \lambda_n \lambda_n + \text{h.c.} \right) + \phi^{*j} (m^2)_j^i \phi_i, \quad (2)$$

where λ_n 's are the Kaluza-Klein modes of the gaugino, and we have assumed a unique gaugino mass M for all λ 's.

We consider the renormalization group (RG) running of the parameters between the fundamental scale $M_{\text{PL}} = M_{\text{Planck}}/\sqrt{8\pi} \simeq 2.4 \times 10^{18}$ GeV and M_{GUT} . To see the gross behavior of the RG running, we consider the contributions coming from only the gauge

supermultiplet, because it is the only source responsible for the power-law running [14,12] of the parameters under the assumptions specified above. We find the following set of the one-loop β functions in this approximation [12,15]:

$$\Lambda \frac{dg}{d\Lambda} = -\frac{2}{16\pi^2} C(G) G_\delta^2 g, \quad (3)$$

$$\Lambda \frac{dM}{d\Lambda} = -\frac{4}{16\pi^2} C(G) G_\delta^2 M, \quad (4)$$

$$\Lambda \frac{dY^{ijk}}{d\Lambda} = -\frac{2}{16\pi^2} (C(i) + C(j) + C(k)) G_\delta^2 Y^{ijk}, \quad (5)$$

$$\Lambda \frac{d\mu^{ij}}{d\Lambda} = -\frac{2}{16\pi^2} (C(i) + C(j)) G_\delta^2 \mu^{ij}, \quad (6)$$

$$\Lambda \frac{dB^{ij}}{d\Lambda} = \frac{2}{16\pi^2} (C(i) + C(j)) G_\delta^2 (2M\mu^{ij} - B^{ij}), \quad (7)$$

$$\Lambda \frac{dh^{ijk}}{d\Lambda} = \frac{2}{16\pi^2} (C(i) + C(j) + C(k)) G_\delta^2 (2MY^{ijk} - h^{ijk}), \quad (8)$$

$$\Lambda \frac{d(m^2)_j^i}{d\Lambda} = -\frac{8}{16\pi^2} C(i) \delta_j^i G_\delta^2 |M|^2, \quad (9)$$

where $G_\delta = gX_\delta^{1/2}(R\Lambda)^{\delta/2}$, and $X_\delta = \pi^{\delta/2}\Gamma^{-1}(1 + \delta/2) = 2(\pi)$ for $\delta = 1(2)$ [12]. The gauge coupling is denoted by g , and $C(G)$ stands for the quadratic Casimir of the adjoint representation of the gauge group G , and $C(i)$ for that of the representation R_i . It is easy to show that the evolution of Y^{ijk} , μ^{ij} and M are related to that of g as

$$M(M_{\text{GUT}}) = \left(\frac{g(M_{\text{GUT}})}{g(M_{\text{PL}})} \right)^2 M(M_{\text{PL}}), \quad (10)$$

$$Y^{ijk}(M_{\text{GUT}}) = \left(\frac{g(M_{\text{GUT}})}{g(M_{\text{PL}})} \right)^{\eta_Y^{ijk}} Y^{ijk}(M_{\text{PL}}), \quad (11)$$

$$\mu^{ij}(M_{\text{GUT}}) = \left(\frac{g(M_{\text{GUT}})}{g(M_{\text{PL}})} \right)^{\eta_\mu^{ij}} \mu^{ij}(M_{\text{PL}}), \quad (12)$$

where the exponents are given as $\eta_Y^{ijk} = (C(i) + C(j) + C(k))/C(G)$ and $\eta_\mu^{ij} = (C(i) + C(j))/C(G)$. Therefore these parameters can become very large if $g(M_{\text{PL}})/g(M_{\text{GUT}})$ is large. A rough estimate shows that

$$\frac{g(M_{\text{GUT}})}{g(M_{\text{PL}})} \simeq \left[\frac{C(G)X_\delta\alpha_{\text{GUT}}}{\pi\delta} \right]^{1/2} \left(\frac{M_{\text{PL}}}{M_{\text{GUT}}} \right)^{\delta/2}. \quad (13)$$

If we use $\alpha_{\text{GUT}} = 0.04$, $M_{\text{PL}}/M_{\text{GUT}} = 10^2$, $G = SU(5)$, then this is given approximately by 3.5 for $\delta = 1$ and 32 for $\delta = 2$.

The ratios between other SSB parameters to the gaugino mass M approach to their infrared attractive fixed points;

$$\begin{aligned} B^{ij}/M &\rightarrow -\eta_\mu^{ij} \mu^{ij}, \\ h^{ijk}/M &\rightarrow -\eta_Y^{ijk} Y^{ijk}, \\ (m^2)_j^i/|M|^2 &\rightarrow (C(i)/C(G)) \delta_j^i, \end{aligned} \quad (14)$$

where η 's are used in (11) and (12). Note that so far no assumption on the reality of the parameters has been made. We see that the low-energy structure is completely fixed by the group theoretic structure of the model. Furthermore, since h^{ijk} and $(m^2)_j^i$ become aligned in the infrared limit, *i.e.*, $h^{ijk} \propto Y^{ijk}$ and $(m^2)_j^i \propto \delta_j^i$, the infrared forms (14) give desired initial values of the parameters at M_{GUT} to suppress FCNC processes in the MSSM, and they predict that the only CP-violating phase is the usual CKM phase.

One can easily estimate how much of a disorder in the initial values at M_{PL} can survive at M_{GUT} . Suppose that there exists an $O(1)$ disorder in $(m^2)_j^i/|M|^2$. Using the β functions (4) and (9), we find the deviation from (14) to be

$$\left(\frac{g(M_{\text{PL}})}{g(M_{\text{GUT}})}\right)^4 \left[\frac{(m^2)_j^i}{|M|^2}(M_{\text{PL}}) - \frac{C(i)}{C(G)}\delta_j^i \right]. \quad (15)$$

Then inserting the value of $g(M_{\text{PL}})/g(M_{\text{GUT}})$ given in (13), we find that an $O(1)$ disorder at M_{PL} becomes a disorder of $O(10^{-2})$ and $O(10^{-6})$ at M_{GUT} for $\delta = 1$ and 2, respectively. Note that the off-diagonal elements of $(m^2)_j^i$ as well as the differences among the diagonal elements $\Delta m^2(i, j) = (m^2)_i^i - (m^2)_j^j$ (if $C(i) = C(j)$) belong to the disorder. However, their contributions to $(\delta_{ij})_{LL, RR}$ of [2] are less than $O(10^{-6})$ for $\delta = 2$, therefore the most stringent constraint coming from $\mu \rightarrow e\gamma$ is satisfied [2]. However, in the case of five dimensions ($\delta = 1$) the suppression of the disorder will not be sufficient. [If we use $M_{\text{PL}}/M_{\text{GUT}} \sim 10^3$, then the suppression is much improved and the five dimensional case is also allowed.]

Similarly, using (4) and (8), we obtain the deviation for the tri-linear couplings from (14) as

$$\left(\frac{g(M_{\text{PL}})}{g(M_{\text{GUT}})}\right)^2 \left[\frac{h^{ijk}}{M}(M_{\text{PL}}) + \eta_Y^{ijk} Y^{ijk}(M_{\text{PL}}) \right], \quad (16)$$

where use has been made of (11). Suppose the tri-linear couplings to be order of MY^{ijk} at M_{PL} . Then we find that

$$\left| \frac{h^{ijk}}{\eta_Y^{ijk} MY^{ijk}}(M_{\text{GUT}}) + 1 \right| \lesssim \left(\frac{g(M_{\text{PL}})}{g(M_{\text{GUT}})} \right)^{2+\eta_Y^{ijk}}. \quad (17)$$

Note that the phases of h^{ijk}/MY^{ijk} can be suppressed. In the case of $G = SU(5)$, $\eta_Y^{ijk} = 48/25(42/25)$ for the up (down) type Yukawa couplings. Using (13) again, we find that the right-hand side of (17) is $\sim 10^{-2(6)}$ for $\delta = 1(2)$. This disorder contributes, for instance, to $Im(\delta_{ii})_{LR}$ as well as $Re(\delta_{ij})_{LR}$ of [2]. Therefore our suppression mechanism can satisfy the most stringent constraints coming from the electric dipole moments (EDM) of the neutron and the electron [2]. Similarly the phases of the B-parameter, $B^{ij}/M\mu^{ij}$ are also suppressed.

In concrete examples, there will be logarithmic corrections to (14) to which the Yukawa couplings Y^{ijk} non-trivially contribute. How much the logarithmic corrections can amplify the disorder will be model-dependent. It is certainly worthwhile to note that the logarithmic interactions will be non-negligible only for Λ close to M_{GUT} , thereby overcoming the problem found in [16] that the GUT effects may destroy the universality of the SSB terms.

To be more specific, we consider the minimal GUT model based on $G = SU(5)$ in six dimensions. The boundary fields are: chiral superfields $\Psi^i(\mathbf{10})$ and $\Phi^i(\bar{\mathbf{5}})$, where i runs over

the three generations, $\Sigma(\mathbf{24})$ to break $SU(5)$, and two Higgs superfields $H(\mathbf{5})$ and $\bar{H}(\bar{\mathbf{5}})$. To simplify the situation, we neglect the neutrino masses and their mixings. The boundary superpotential of the model is given by

$$W = \frac{Y_U^{ij}}{4} \Psi^i \Psi^j H + \sqrt{2} Y_D^{ij} \Phi^i \Psi^j \bar{H} + \frac{Y_\lambda}{3} \Sigma^3 + Y_f \bar{H} \Sigma H + \frac{\mu_\Sigma}{2} \Sigma^2 + \mu_H \bar{H} H, \quad (18)$$

where Y_U^{ij} and Y_D^{ij} are the Yukawa couplings. Then the gross infrared attractive form of the SSB parameters (14) becomes

$$B_\Sigma \rightarrow -2M\mu_\Sigma, \quad B_H \rightarrow -\frac{24}{25}M\mu_H, \quad (19)$$

$$h_U^{ij} \rightarrow -\frac{48}{25}MY_U^{ij}, \quad h_D^{ij} \rightarrow -\frac{42}{25}MY_D^{ij}, \quad (20)$$

$$(m_\Phi^2)^{ij} \rightarrow \frac{12}{25}|M|^2\delta^{ij}, \quad (m_\Psi^2)^{ij} \rightarrow \frac{18}{25}|M|^2\delta^{ij}, \quad (21)$$

$$m_{H_d}^2, m_{H_u}^2 \rightarrow \frac{12}{25}|M|^2, \quad (22)$$

in an obvious notation. We find also $h_f \rightarrow -(49/25)MY_f$, $h_\lambda \rightarrow -3MY_\lambda$, $m_\Sigma^2 \rightarrow 2|M|^2$. All the scalars that belong to $\mathbf{5}$ or $\bar{\mathbf{5}}$ have the same positive squared soft-mass ($\approx (0.69M)^2$), which does not differ very much from that ($\approx (0.85M)^2$) for the scalars belonging to $\mathbf{10}$. So, the infrared attractive form in the present model is similar to the SSB terms of the constrained MSSM (CMSSM), and therefore, the model predicts a similar spectrum as in the CMSSM.

In the following analyses we would like to neglect the mixings of the matter multiplets, because their effects will be very small as seen later. One of the pleasant feature of the infrared attractive form of the SSB terms (14) is that the tri-linear couplings, too, may be assumed to be small if the corresponding Yukawa couplings are small, as we have seen in (16). Consequently, we will approximate $Y_{U,D}^{ij}$ by $Y_{t,b}\delta^{i3}\delta^{j3}$ and $h_{U,D}^{ij} = h_{t,b}\delta^{i3}\delta^{j3}$, respectively.

To proceed, we require that the MSSM is the effective theory below M_{GUT} , and we would like to check whether the initial values of the SSB terms at M_{GUT} given in (19)–(21) cause conflicts at low-energies, especially with electroweak symmetry breaking. For simplicity, we assume that the potential of the MSSM at M_{SUSY} takes the tree-level form, where we identify M_{SUSY} with the unified gaugino mass M . We require that $M_t = 174$ GeV and $M_\tau = 1.77$ GeV, while imposing the $b - \tau$ unification at M_{GUT} . [But we will not take the mass of the bottom quark very seriously. It becomes slightly larger than its experimental value.]

We are also interested in how much the Yukawa interactions modify the infrared attractive values (19)–(21). Because of the limitation of space, we present below only one case, which is consistent with electroweak symmetry breaking. The parameters of the case are: $M = 500$ GeV, $g = (0.0406 \times 4\pi)^{1/2}$, $M_{\text{GUT}} = 1.83 \times 10^{16}$ GeV, and

$$\mu_H = 935 \text{ GeV}, \quad Y_t = 0.767g, \quad Y_b = 0.201g, \quad Y_f = 1.0g, \quad Y_\lambda = 0.01g, \quad (23)$$

where $\tan\beta$ is found to be 19.5. In this case the infrared attractive values of the SSB terms are given by

$$\begin{aligned}
(m_{\Phi^{1,2}}^2, m_{\Phi^3}^2) &= (0.534, 0.531)|M|^2, \quad (m_{\Psi^{1,2}}^2, m_{\Psi^3}^2) = (0.801, 0.759)|M|^2, \\
(m_{H_u}^2, m_{H_d}^2) &= (0.383, 0.420)|M|^2, \\
h_t &= -1.97MY_t, \quad h_b = -1.74MY_b, \quad B_H = -0.922M\mu_H.
\end{aligned} \tag{24}$$

We have used the notation $m_{\Phi^i}^2 = (m_{\Phi}^2)^{ii}$, and similarly for $m_{\Psi^i}^2$. It should be noted also that any charged sparticle does not become a LSP with these soft scalar masses at M_{GUT} .

The β functions for $m_{\Phi^{1,2}}^2$ and $m_{\Psi^{1,2}}^2$ do not depend on Y_i and h_i ($i = t, b, f, \lambda$) in our approximation. Therefore, the infrared attractive values (21) are not modified by them. There exist of course logarithmic corrections coming from the gauge interaction, but they are flavor-blind. This is very pleasant, because the most stringent constraint from FCNC processes is the almost degeneracy of the squared soft masses of the first two generations. We have found that for the initial values of Y 's and g given in (23), the off-diagonal components $(m_{\Phi}^2)^{ij}/|M|^2$ and the difference of diagonal elements, $\Delta m_{\Phi}^2(1, 2)/|M|^2 = |m_{\Phi^1}^2 - m_{\Phi^2}^2|/|M|^2$ (and similarly for m_{Ψ}^2) are less than $O(10^{-4})$, which has been estimated to be $O(10^{-6})$ without the logarithmic corrections in (15). This order of disorder at M_{GUT} is still sufficient to satisfy the most stringent constraint coming from $\mu \rightarrow e\gamma$ [2]. In contrast to the case of the first two generations, the β functions for $m_{\Phi^3}^2$ and $m_{\Psi^3}^2$ depend on Y_i and h_i . Therefore, they change their infrared attractive values. Fig. 1 shows the evolution of $m_{\Phi}^2/|M|^2$ and $m_{\Psi}^2/|M|^2$, respectively. The dashed lines correspond to the third generation. We find that the differences $\Delta m_{\Psi}^2(i, 3)/|M|^2 = |m_{\Psi^i}^2 - m_{\Psi^3}^2|/|M|^2$ with $i = 1, 2$ at M_{GUT} are $\lesssim 0.04$, and they contribute to $\mu \rightarrow e\gamma$ through the mixing of the first two and the third generations in the lepton sector, *i.e.*, $V_{3i}^{e*}V_{3j}^e$ [2]. Assuming that V^e can be approximated by the usual Cabibbo-Kobayashi-Maskawa matrix V_{CKM} , we find that $(\delta_{12}^l)_{RR} \lesssim 2 \times 10^{-5}$ in the present case, so that $\Delta m_{\Psi}^2(i, 3)/|M|^2$ does not cause a problem with $\mu \rightarrow e\gamma$ [2]. Similarly, the universality between $m_{H_d}^2$ and $m_{H_u}^2$ are also destroyed, as we see in (24). The main origin are the top Yukawa coupling Y_t and Y_f . This does not conflict with the FCNC problems and CP-violating processes.

In Fig. 2 the converging behavior for $-h_t/MY_t$ and $-h_b/MY_b$ are presented. There is also no universality between h_U and h_D from the beginning. In (16) we have found that the non-aligned part of h^{ijk} is suppressed by a factor of 10^{-6} in six dimensions, if the Yukawa couplings are neglected. Let us estimate how much of this suppression can survive if Y 's are taken into account. We find that the corrections can be written as $\Delta h_U^{ij}/M \sim (1/16\pi^2)(a_t Y_U^{i3} Y_U^{3j} Y_t + a_b Y_U^{i3} Y_D^{3j} Y_b) \ln \Lambda_{\text{eff}}/M_{\text{GUT}}$ and similarly for $\Delta h_D^{ij}/M$, where a_t and a_b are $O(1)$ constants, and we have assumed that $h_{U,D}$ are proportional to $MY_{U,D}$ at a scale Λ_{eff} , at which Y 's become non-negligible. Further considerations in the basis where Y_U is diagonal yield that nonzero contributions (that are relevant to us) are: $|\Delta h_U^{3j}(j \neq 3)| \sim Y_t Y_b^2 L$, $|\Delta h_D^{i3}(i \neq 3)| \sim V_{CKM}^{ib} Y_t^2 Y_b L$, $|\Delta h_D^{ij}(i, j \neq 3)| \sim V_{CKM}^{ib} Y_b^3 L$, where $L = M \ln(\Lambda_{\text{eff}}/M_{\text{GUT}})/16\pi^2$. Assuming that $\Lambda_{\text{eff}} \sim 50M_{\text{GUT}}$, we find that $|\Delta h_U^{3j}(j \neq 3)|/M \sim O(10^{-4})$ for the values given in (23), and the other Δh 's receive a further suppression from V_{CKM} . $\text{Im}\Delta h_D^{11}$, for instance, contributes to the EDM of the neutron, and can be estimated to be $O(10^{-7})$. Therefore, we may conclude that the disorder of the trilinear couplings caused by the Yukawa couplings are sufficiently suppressed to satisfy even the most stringent constraints from the EDMs [2].

We conclude that gauge interactions in extra dimensions can be used to suppress the disorder of the SSB terms at the fundamental scale so that the FCNC processes and dangerous

CP-violating phases become tiny at lower energy scales. Thus the smallness of FCNC may be a possible hint of the existence of extra dimensions.

FIGURES




FIG. 1. Infrared attractiveness of $m^2_\Phi/|M|^2$ and $m^2_\Psi/|M|^2$. The dashed (solid) lines correspond to the third (first two) generation(s). $m^2_{\Psi_{1,2}} > m^2_{\Psi_3} > m^2_{\Phi_{1,2}} \simeq m^2_{\Phi_3}$ at $\Lambda = M_{\text{GUT}}$.




FIG. 2. Infrared attractiveness of $-h_t/M$ (solid) and $-h_b/M$ (dashed).

REFERENCES

- [1] For a recent review, see, for instance, G. L. Kane, hep-ph/0202185, and references therein.
- [2] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57**, 961 (1986); F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, Nucl. Phys. **B477**, 321 (1996); R. Barbieri, L. Hall and A. Strumia, Nucl. Phys. **B445**, 219 (1995); A. Bartl, T. Gajdosik, E. Lunghi, A. Masiero, W. Porod, H. Stremnitzer and O. Vives, Phys. Rev. **D64**, 076009 (2001), and references therein.
- [3] M. Dine and A. E. Nelson, Phys. Rev. **D48**, 1277 (1993); M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. **D51**, 1362 (1995); For a review, G. F. Giudice and R. Rattazzi, Phys. Rept. **322**, 419 (1999).
- [4] L. Randall and R. Sundrum, Nucl. Phys. **B557**, 79 (1999); G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **12**, 027 (1998).
- [5] D. E. Kaplan, G. D. Kribs and M. Schmaltz, Phys. Rev. **D62**, 035010 (2000); Z. Chacko, M. A. Luty, A. E. Nelson and E. Ponton, JHEP **01**, 003 (2000).
- [6] A. Anisimov, M. Dine, M. Graesser and S. Thomas, hep-th/0111235; hep-th/0201256.
- [7] A. E. Nelson and M. J. Strassler, JHEP **09**, (2000) 030; hep-ph/0006251; T. Kobayashi and H. Terao, Phys. Rev. **D64**, 075003 (2001).
- [8] T. Kobayashi, H. Nakano and H. Terao, Phys. Rev. **D65**, 015006 (2002).
- [9] M. A. Luty and R. Sundrum, Phys. Rev. **D65**, 066004 (2002); hep-th/0111231.
- [10] A. Karch, T. Kobayashi, J. Kubo, and G. Zoupanos, Phys. Lett. **B441**, 235 (1998); M. Luty and R. Rattazzi, JHEP **11**, 001 (1999).
- [11] I. Antoniadis, Phys. Lett. B **246**, 377 (1990); I. Antoniadis, C. Muñoz and M. Quirós, Nucl. Phys. **B397**, 515 (1993).
- [12] K. Dienes, E. Dudas and T. Gherghetta, Nucl. Phys. **B537**, 47 (1999).
- [13] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Rev. **D59**, 086004 (1999).
- [14] T. R. Taylor and G. Veneziano, Phys. Lett. B **212**, 47 (1988).
- [15] T. Kobayashi, J. Kubo, M. Mondragon and G. Zoupanos, Nucl. Phys. **B550**, 99 (1999).
- [16] N. Polonsky and A. Pomarol, Phys. Rev. Lett. **73**, 2292 (1994).